

The Shape and History of the Moon

- Presented to *The Lunar Science Forum*


July 19, 2012

- By Charles J. Byrne

Image Again

charles.byrne@verizon.net

www.imageagain.com



The Dichotic Moon


- The “Man in the Moon” (maria patterns) are mostly on the near side
- The near side is low, a bulge on the far side
- The crust is thin on the near side, thick on the far side
- Heavy element anomalies are mostly on the near side
- Moments of inertia are uneven
- Center of gravity is offset to the near side

-
-
-

Near Side of the Moon



Far Side of the Moon



Maps of the Current Moon


- Topography
 - Photography
 - Digital Elevation Map (DEM)
- Gravity
- Crustal thickness
- Mineral concentrations

Maps of the Current Moon


- Topography
 - Photography
 - Digital Elevation Map (DEM)
- Gravity
- Crustal thickness
- Mineral concentrations

Digital Elevation Map (DEM)


North Pole




Impact Model: Maxwell-Z



Crater Depth vs. Diameter




Scaling by Depth and Diameter



-
-
-


Models of Basins and Large Craters



The Big Guys: Megabasins

- A megabasin is a basin that contains other basins
- A model of a megabasin must consider the spherical nature of the Moon
- Ejecta is thrown into elliptical orbit: velocity as a function of radius is needed
- Near the antipode, ejecta is focused: the depth of the ejecta field increases

Scaled Velocity Profile




The Search for Megabasins

- Model parameters : center latitude and longitude, diameter, depth, and fill
- Two megabasins were modeled together: the South Pole-Aitken Basin (SPA) and a mystery basin
- Parameters were varied to best fit the Moon
- Both the Near Side Megabasin (NSM) and its ejecta field, the far side bulge, emerged.
- The St. John-Teselius Basin emerged from the residual DEM of SPA and NSM.


-
-
-

Three Megabasins




-
-
-

Megabasins, Impacts, and Maria




-
-
-

Interim Residual DEM



•
•
•


Mounds and Depressions



• • • • • • • • • •

-
-
-


Comprehensive Model




-
-
-

Model Compared to Topography



Model




Topography



The NSM and its Antipode




Radial Profile of the NSM




SPA Impact Simulation (Stewart)


A
000




B
500



C
2000



D
4000




History of the Moon


- Accretion 4.5 Ga
- Megabasins 4.34 - ?
- pre-Nectarian Period 4.34 - 4.0 Ga
- Nectarian Period 4.0 - 3.9 Ga
- Early Imbrian Period 3.9 - 3.8 Ga
- Later Imbrian Period 3.8 - 3.2 Ga
- Eratosthenian Period 3.2 - 0.8 Ga
- Copernican Period 0.8 - 0 Ga

-
-
-


Megabasins (4.34 Ga - ?)



Pre-Nectarian (700 Ma)




Nectarian (100 Ma)




-
-
-

Lower Imbrian (100 Ma)




Upper Imbrian (400 Ma)




-
-
-

Upper Imbrian + Maria




Erastosthenian (2400 Ma)



-
-
-

Copernican (800 Ma to present)



Summary

- Major features of the Moon's surface have been deconstructed from topography
- Three megabasins established the general topography and crustal thickness
- Intensive bombardment followed and volcanic lava erupted through thinned crust
- Mineral anomalies are associated with the melt columns of the NSM and SPA

-
-
-

Questions?

